

Bred for Forages Tall & Early



KWS AVIATOR

KWS AVIATOR has been bred for top forage yields and backcrossed with a variety that has great winter hardiness and early season vigor. KWS AVIATOR is expected to provide ruminant livestock producers with an earlier harvested product, and when used in rotation with other forage crops will extend the forage harvest season.

Forage Production Test Data

	KWS AVIATOR	KWS PROGAS	KWS PROPOWER
Early Cut (Flag Leaf – Boot) MT/AC @ 35% DM	7.22	6.56	6.17
Late Cut (Milky – Soft Dough) MT/AC @ 35% DM	12.04	11.20	11.62

Source: Third Party USA Rye Trials
Kansas City, KS | Hayes, KS | Scandia, KS | Cornell U, NY | SDSU, SD | 2021-2022

Variety Highlights

- Top forage yields
- Tall biomass structure
- Earliest spring emergence
- Unsurpassed winter hardiness

Agronomy Co-op Test Data

	KWS AVIATOR	Average of Hybrid Rye Check Varieties
Winter Hardiness	91%	81%
Height	103 cm	92 cm

Source: Fall Rye Cooperative Registration Trials 2016-17, 2017-18, 2018-19

For more information contact us:

@ canada@kws.com

☎ 403.926.8861



Scan to watch our KWS PollenPLUS video

KWS RyeGHT Seeding Practices

Forage Production | Western Canada

Hybrid fall rye can be an easy crop to grow and success of growing is primarily achieved from having strong stand establishment in the fall. These practices will optimize both performance and consistency year after year, starting with the most important first.

RyeGHT Time:

General seeding period for hybrid fall rye in Western Canada is from August 15 to September 25. It has consistently been shown that **early seeding** provides hybrid fall rye the best opportunity to achieve maximum root and tiller growth for the highest yield and quality.

RyeGHT Rate:

The standard seeding rate is **1 unit/ac** under normal conditions.

- August 15-31: Seed at 1 to 1.2 units/ac into a firm seedbed to achieve 5-6" shoot growth with 6-8 tillers/plant. Under irrigation 0.8 units/ac may be used.
- September 1-15: Seed at 1.2 units/ac into a firm seedbed to achieve 3-5" of shoot growth with 4-5 tillers/plant. Under irrigation 0.8 to 1 units/ac may be used.
- September 15-25: Seed at 1.2 to 1.3 units/ac into a firm seedbed to achieve 2-3" of shoot growth with 2-3 tillers/plant. Under irrigation 1 to 1.2 units/ac may be used.

RyeGHT Depth:

Always seed shallow at **¾" depth** with travel speed <4.5 mph to maintain consistent seed depth. Seeding deeper or faster will result in higher seedling mortality and reduced stand. Maintain **¾"** seed depth even if soil is dry and wait for rain.

RyeGHT Stand:

The plant stand target is **16-20 plants/ft²**. Even emergence and achieving a target stand will provide the most consistent yields.

RyeGHT Field Selection:

Hybrid rye grows best on well drained fields, free of residual herbicides that affect rye. It is the first choice to grow on land with stubble cover. It can also be grown on land without stubble provided it is seeded early, maximizes shoot growth, and tillers per plant. Areas prone to wind erosion should be seeded at a higher rate.

RyeGHT Trash Management:

It is important to spread previous crop straw and chaff evenly and as close to the full width of the header as possible. Harrowing can be beneficial; however, it can also dislodge residual plants from the previous crop causing residue piles and seed placement issues. Use higher seeding rates when seed bed issues exist.

